

ALGORITHMIC REDUCTION AND RATIONAL GENERAL SOLUTIONS OF FIRST ORDER ALGEBRAIC DIFFERENTIAL EQUATIONS

GUOTING CHEN AND YUJIE MA

ABSTRACT. First order algebraic differential equations are considered. An necessary condition for a first order algebraic differential equation to have a rational general solution is given: the algebraic genus of the equation should be zero. Combining with Fuchs' conditions for algebraic differential equations without movable critical point, an algorithm is given for the computation of rational general solutions of these equations if they exist under the assumption that a rational parametrization is provided. It is based on an algorithmic reduction of first order algebraic differential equations with algebraic genus zero and without movable critical point to classical Riccati equations.

1. INTRODUCTION

The study of first order algebraic differential equations can be dated back to C. Briot and T. Bouquet [2], L. Fuchs [11] and H. Poincaré [20]. M. Matsuda [17] gave a modern interpretation of the results using the theory of differential algebraic function field of one variable, and Eremenko [9] studied the bound of the degrees of the rational solutions of a first order algebraic differential equation by using the approach of [17].

From an algorithmic point of view, many authors have been interested in the constructions of closed form solutions for differential equations (this problem can be traced back to the work of Liouville). In [21], Risch, gave an algorithm for finding closed forms for integration. In [14], Kovacic presented a method for solving second order linear homogeneous differential equations. In [26], Singer proposed a method for finding liouvillian solutions of general linear differential equations. In [15], Li and Schwarz gave a method to find rational solutions for a class of partial differential equations. All these works are limited to linear cases.

For algebraic (nonlinear) differential equations there are some studies in this direction. For Riccati equations, polynomial solutions are considered in [4] and algorithms for the computation of rational solutions are given in [3, 14]. In [5], the algebraic solutions of a general class of first order and first degree algebraic differential equations was studied and the degree bound of algebraic solutions in the nondicritical case was given. In [10, 16], algorithms for the computation of rational

Date: January 1, 2004.

1991 Mathematics Subject Classification. Primary 34A09, 68W30; Secondary 14Q05, 34M15.

Key words and phrases. Algebraic differential equation, rational general solution, rational parametrization.

The second author is partially supported by the NKBRSF of China (No. 2004CB318000), the NNSF (No. 10301032) and by a CNRS—K. C. WONG fellowship during his visit to the Laboratoire P. Painlevé, Université de Lille 1, France.

general solutions or polynomial solutions are given for some kinds of algebraic differential equations.

Another motivation of our work is differential algebraic geometry. In the series papers of Wu [29, 30, 31], the author studied algebraic differential geometry from several different points of view. In [31], the author presents an algorithmic method of solving arbitrary systems of algebraic differential equations by extending the characteristic set method to differential case. The Devil's problem of Pommaret is solved in details as an illustration.

In this paper, we consider the computation of rational general solutions of first order algebraic differential equations by using methods from algebraic geometry. We give an necessary condition for a first order algebraic differential equation to have a rational general solution: that is the algebraic genus of the equation should be zero. Combining with Fuchs' conditions for first order algebraic differential equations without movable critical points, we obtain an algorithm for the computation of rational general solutions under the assumption that a rational parametrization is provided. It is based on an algorithmic reduction of first order algebraic differential equation of algebraic genus zero and without movable critical point to classical Riccati equations.

2. RATIONAL GENERAL SOLUTIONS OF FIRST ORDER ALGEBRAIC DIFFERENTIAL EQUATIONS

We first present some results from algebraic geometry, which is used in the following.

Let $f(x, y)$ be an irreducible polynomial over $\mathbb{C}$. We call that $f(x, y) = 0$ is a rational curve if there exist two rational functions $\phi(t), \psi(t) \in \mathbb{C}(t)$ such that

- (i) For all but a finite set of $t_0 \in \mathbb{C}$, $(\phi(t), \psi(t))$ is a point of f .
- (ii) With a finite number of exceptions, for every point (x_0, y_0) of f there is a unique $t_0 \in \mathbb{C}$ such that $x_0 = \phi(t_0)$, $y_0 = \psi(t_0)$.

It is impossible to avoid having a finite number of exceptions in the above conditions. They arise from two sources. One is the fact that a rational function is not defined for some values of the variable, and the other is the presence of singular points on the curve.

The following results are well known in algebraic geometry [28].

Theorem 2.1. *An algebraic curve is rational if and only if its genus is zero.*

Theorem 2.2. (a) *Every rational transform of a rational curve is a rational curve.*
 (b) *If λ is transcendental over $\mathbb{C}$ and if $\mathbb{C} \subset F \subset \mathbb{C}(\lambda)$, $F \neq \mathbb{C}$, then there is an element $\mu \in F$, transcendental over $\mathbb{C}$, such that $F = \mathbb{C}(\mu)$.*
 (c) *If a curve $f(x, y) = 0$ satisfies (i) for rational functions $\phi(\lambda), \psi(\lambda)$ which are not both constants, then there exist rational functions $\tilde{\phi}(\lambda), \tilde{\psi}(\lambda)$ for which both (i) and (ii) are satisfied, and the curve is rational.*

The three statements in Theorem 2.2 are all equivalent. It is called Lüroth's theorem.

Now we consider a first order algebraic differential equation in the form

$$(2.1) \quad F(z, w, w') = 0,$$

where F is a polynomial in w, w' with rational coefficients in $\mathbb{C}(z)$.

Definition 2.3. A general solution $w(z, \lambda)$ of (2.1) is called a rational general solution if it is rational in z and λ .

If $F(z, w, w') = 0$ admits a rational general solution, then it is free from movable critical point for poles are the only singularities of the solution which change their position if one varies the initial data $c \in \mathbb{C}$.

We now prove the following theorem on rational general solution of a first order algebraic differential equation.

Theorem 2.4. *If a first order irreducible algebraic differential equation*

$$F(z, w, w') = 0$$

admits a non-constant rational general solution, then the genus of $F(z, w, w') = 0$ with respect to w, w' is zero for any z , except for a finitely many exceptions.

Proof. Let $w = r(z, \lambda)$ be the rational general solution of $F = 0$ with the arbitrary constant λ . Then $w(z) = r(z, \lambda)$ and $w'(z) = \frac{\partial r}{\partial z}$ are rational functions and they satisfy equation $F(z, w, w') = 0$.

Let z be fixed and consider the curve $f_z(x, y) = F(z, x, y) = 0$. Denote $\phi_z(\lambda) = w(z)$ and $\psi_z(\lambda) = w'(z)$. If $\psi_z(\lambda)$ is a constant, then $w(z) = zw'(z) + \lambda$ is of genus zero and hence the genus of $f_z(x, y) = 0$ is also zero. If $\psi_z(\lambda)$ is not a constant, consider the point $(\phi_z(\lambda), \psi_z(\lambda))$ of $f_z(x, y) = 0$ for the parameter λ in the transcendental extension field $\mathbb{C}(\lambda)$. It is clear that (i) is satisfied for all but finitely many λ . Hence $f_z(x, y) = 0$ is a rational curve and its genus is zero by Theorem 2.1 and Theorem 2.2. $\square$

Motivated by this theorem, we present the following definition.

Definition 2.5. The algebraic genus of a first order algebraic differential equation $F(z, w, w') = 0$ is defined to be the genus of $F(z, w, w') = 0$ with respect to w and w' .

3. REDUCTION OF FIRST ORDER ALGEBRAIC DIFFERENTIAL EQUATIONS

For a first order algebraic differential equation $F(z, w, w') = 0$, Fuchs Theorem presented necessary conditions for the equation to be free from movable critical point. By the Painlevé Theorem, we know that Fuchs' conditions are sufficient (see [11, 13, 18, 20]).

3.1. Fuchs Theorem. Let $D(z, w)$ be the p -discriminant of the equation $F(z, w, w') = 0$, it is a polynomial in w , whose coefficients are analytic functions of z [13].

The conditions, necessary to secure that the first order differential equation

$$F(z, w, w') = 0$$

of degree m , shall have no movable critical point, are:

- (1) The coefficient $A_0(z, w)$ is independent of w and therefore reduces to a function of z alone or to a constant. The equation may then be divided throughout by A_0 and takes the form

$$w'^m + \psi_1(z, w)w'^{m-1} + \cdots + \psi_{m-1}(z, w)w' + \psi_m(z, w) = 0$$

in which the coefficients ψ are polynomials in w , and analytic, except for isolated singular points, in z .

- (2) If $w = \eta(z)$ is a root of $D(z, w) = 0$, and $p = \omega(z)$ is a multiple root of $F(z, \eta, \eta') = 0$, such that the corresponding root of $F(z, w, w') = 0$, regarded as a function of $w - \eta(z)$ is branched, then

$$\omega(z) = \frac{d\eta}{dz}.$$

- (3) If the order of any branch is α , so that the equation is effectively of the form

$$\frac{d}{dz}\{w - \eta(z)\} = c_k\{w - \eta(z)\}^{\frac{k}{\alpha}}$$

then $k \geq \alpha - 1$.

3.2. Reduction to Classical Riccati Equation. Consider now a first order algebraic differential equation $F(z, w, w') = 0$ of genus zero and without movable critical point. One can find a parametrization of the rational curve $F(z, x, y) = 0$ in the form $x = r_1(t, z)$ and $y = r_2(t, z)$ with $r_1(t, z)$ and $r_2(t, z)$ rational functions in t and z . By the inversion of rational curves we know that t is rational function in z, x and y . For algorithm on parametrization and inversion of rational curves we refer to [1, 22, 23, 24, 25, 27]. One has

$$(3.1) \quad \frac{dt}{dz} = \left(r_2(t, z) - \frac{\partial r_1}{\partial z} \right) / \frac{\partial r_1}{\partial t} = \frac{P(t, z)}{Q(t, z)},$$

where P and Q are polynomials in t and z . Since $F(z, w, w') = 0$ has no movable critical point, one knows that (3.1) is also has no movable critical point as t is rational function in z, x and y . By Fuchs Theorem, we have that equation (3.1) is a Riccati equation ([12] Chapter II, §7), that is

$$(3.2) \quad \frac{dt}{dz} = A(z)t^2 + B(z)t + C(z),$$

where A, B, C are rational functions in z . We distinguish two cases according to $A(z)$.

Case 1: If $A(z) \not\equiv 0$, we consider the change of variables $t(z) = -u(z)/A(z)$. One has

$$u'(z) + u^2 = (B(z) + A'(z)/A(z))u - C(z)A(z),$$

In which the coefficient $A(z)$ is reduced to -1 . Next we make the change $u = v + \beta(z)$ to reduce the coefficient of u to zero by choosing an appropriate β . We obtain finally a classical Riccati equation in the form

$$(3.3) \quad v' + v^2 = r(z) \in \mathbb{C}(z).$$

Algorithms for the computations of rational solutions of classical Riccati equations are given in many literatures (see for example [3, 14]).

If $r(z) \not\equiv 0$ then a rational solution of equation (3.3) is equivalent to an exponential solution $e^{\int v(z)dz}$ of the linear differential equation

$$(3.4) \quad y'' = r(z)y.$$

Proposition 3.1. *If the Riccati equation (3.3) with $r(z) \not\equiv 0$ has a general rational solution, then $r(z)$ has the form*

$$r(z) = \sum_{i=1}^m \left(\frac{\beta_i}{(z - z_i)^2} + \frac{\gamma_i}{(z - z_i)} \right),$$

in which $4\beta_i = n_i^2 - 1$ where n_i is an integer ≥ 2 .

Proof. Suppose that $v(z)$ is a rational solution of equation (3.3). Let $z_1, \dots, z_m$ be the poles of r . According to Kovacic's algorithm (see [14]), $v(z)$ should be in the form

$$v(z) = \sum_{i=1}^m \sum_{j=1}^{\nu_i} \frac{a_{ij}}{(z - z_i)^j} + \sum_{k=1}^d \frac{1}{z - c_k} + f(z),$$

where the ν_i are known, a_{ij} are known up to a two choices each, d is known, and $f \in \mathbb{C}[z]$ is known up to two choices. Hence there may be arbitrary parameter only in the determination of the c_k .

Let

$$P(z) = \prod_{k=1}^d (z - c_k),$$

and

$$\omega(z) = \sum_{i=1}^m \sum_{j=1}^{\nu_i} \frac{a_{ij}}{(z - z_i)^j} + f(z).$$

Then $v = P'/P + \omega$ and $y = e^{\int v} = P e^{\int \omega}$ is a solution of the linear differential equation (3.4). Hence P is a polynomial solutions of degree d of the following linear equation:

$$(3.5) \quad P'' + 2\omega P' + (\omega' + \omega^2 - r)P = 0.$$

One can determine whether it has a general polynomial solution or not.

Furthermore if (3.3) admits a rational general solution, then writing $r(z) = p(z)/q(z)$, according to [32], one has

- (a) $\deg(p) - \deg(q) \leq -2$;
- (b) $r(z)$ has only double poles, hence

$$r(z) = \sum_{i=1}^m \left(\frac{\beta_i}{(z - z_i)^2} + \frac{\gamma_i}{(z - z_i)} \right),$$

in which $4\beta_i = n_i^2 - 1$ where n_i is an integer ≥ 2 .

□

Therefore a possible rational solution of equation (3.3) with $r(z)$ as in the proposition should be

$$v(z) = \sum_{i=1}^m \frac{a_i}{z - z_i} + \sum_{k=1}^d \frac{1}{z - c_k},$$

where $a_i^2 - a_i - \beta_i = 0$. Hence to determine a rational general solution one needs to compute polynomial solutions of equation (3.5) in order to determine the c_k .

Case 2: If $A(z) \equiv 0$, then one can integrate easily the linear equation

$$t' = B(z)t + C(z)$$

to get the general solution

$$t(z) = \left(\int C(z)dz + \lambda \right) e^{\int B(z)dz},$$

where λ is an arbitrary constant. Effective algorithm is given in [21] for integration in closed forms. One may find rational solutions in this way. It is clear that in this

case one may get rational general solutions only if both $\int C(z)dz$ and $e^{\int B(z)dz}$ are rational functions.

3.3. First Order Algebraic Differential Equations with Constant Coefficients. As an application, we consider a first order algebraic differential equation with constant coefficients,

$$(3.6) \quad F(w, w') = 0.$$

This kind of equations was systematically studied in [2] and an algorithm is given in [10] for the determination of the rational general solution of the equation if it exists.

When using the above reduction for equation (3.6), it is clear that z is not involved in the equation, we henceforth get a Riccati equation with constant coefficients.

As above there are two cases to be considered. In case 1, if equation (3.6) has a non constant rational solution $w(z)$ then $w(z + \lambda)$ is a general rational solution. Since the equation $u' + u^2 = c$ for a constant $c \neq 0$ does not have a rational solution, then one can reduce equation (3.6) to the equation $u' + u^2 = 0$ in which case we have the general solution

$$u = \frac{1}{z + \lambda}.$$

In case 2, the equation can be converted to $u' = bu + c$ with constant b, c . Then $u = \lambda e^{bz} - \frac{c}{b}$ if $b \neq 0$, and $u = cz + \lambda$ if $b = 0$, where λ is an arbitrary constant.

Summarizing the above, we then have the following

Corollary 3.2. *Let $F(w, w') = 0$ be a first order irreducible algebraic differential equation with constant coefficients. Then it has a non-constant rational general solution if and only if it can be reduced either to a linear equation $u' = c$ for some constant c or to a Riccati equation of the form $u' + u^2 = 0$.*

Example 1. Consider

$$F(y, y') = y'^4 - 8y'^3 + (6 + 24y)y'^2 + 257 + 528y^2 - 256y^3 - 552y.$$

This example comes from [10]. One finds by computation in Maple that its algebraic genus is zero and it has the following rational parametrization.

$$y = \frac{17}{16} - 27t + \frac{2187}{2}t^2 + 531441t^4, \quad y' = 78732t^3 + 81t - 1.$$

And the corresponding Riccati equation is $t'(z) = \frac{1}{27}$. Hence $t = \frac{1}{27}z + \lambda$ and the general solution of the differential equation $F(y, y') = 0$ is

$$y(z) = \frac{17}{16} - 27\left(\frac{1}{27}z + \lambda\right) + \frac{2187}{2}\left(\frac{1}{27}z + \lambda\right)^2 + 531441\left(\frac{1}{27}z + \lambda\right)^4,$$

where λ is an arbitrary constant.

4. ALGORITHM AND EXAMPLE

We can now give the following algorithm on seeking for a rational general solution of a first order algebraic differential equation.

4.1. Algorithm. Algorithm:

Input: $F(z, w, w') = 0$ is a first order algebraic differential equation.

Output: A rational general solution of $F(z, w, w') = 0$ if it exists.

- (1) Determine the irreducibility of the equation.
If $F(z, w, w') = 0$ is reducible, then factorize it and go to step (2) for each branch curve of $F(z, w, w') = 0$, else go to step 2 directly.
- (2) Compute the algebraic genus g of $F(z, w, w') = 0$. If $g \neq 0$, then the equation doesn't admit rational general solution by Theorem 2.4, else go to step 3.
- (3) Determine the Fuchs' conditions of $F(z, w, w') = 0$. If the conditions are not satisfied, the algorithm terminates, else go to step 4.
- (4) If a rational parametrization of $F(z, w, w') = 0$ in the form $w = r_1(t, z)$ and $w' = r_2(t, z)$ with $r_1(t, z)$ and $r_2(t, z)$ rational functions in t and z is provided, go to step 5.
- (5) Compute the derivative $\frac{dt}{dz} = (r_2(t, z) - \frac{\partial r_1}{\partial z}) / \frac{\partial r_1}{\partial t}$ which is a Riccati equation of the form (3.2) by the Fuchs Theorem.
- (6) Reduce the above Riccati equation to a classical Riccati equation (3.3) and compute a rational solution using the algorithm in [3, 14].

Remark 4.1. Assume that $F(z, w, w') = 0$ has algebraic genus 0 for all z (or for all z with some finite exceptions), in general, the rational parametrization that we obtain using the the algorithm in [1, 24, 25, 27] are not rational functions in z (it could be appear algebraic elements over $\mathbb{C}(z)$ as $\sqrt{z}$ (see [25])). Our algorithm works only with for those equation $F(z, w, w') = 0$ for which a rational parametrization is provided. For instance, for first order algebraic differential equations with constant coefficients.

Remark 4.2. Our algorithm for the case of constant coefficients is the same as the classical one, which is for instance described in [19], Ch. IV, sec. I (page 62–65).

4.2. Example. Example 2. Consider the following equation:

$$F(z, w, w') = w'^2 + \frac{2w}{z}w' - 4zw^3 + \frac{(1 + 12z^2)w^2}{z^2} - 12\frac{w}{z} + \frac{4}{z^2}.$$

Its algebraic genus is zero. One gets the following rational parametrization by Maple:

$$\begin{aligned} w = r_1 &= \frac{t^2z^2 + 4t^2 - 6tz + 1 + 4z^2}{4z(-z + t)^2}, \\ w' = r_2 &= -\frac{-4z^3 + 13tz^2 + t + 2t^2z^3 - 10t^2z + t^3z^4 + t^3z^2 + 4t^3}{4z^2(-z + t)^3}. \end{aligned}$$

One obtains the following Riccati equation

$$t' = \frac{(z^2 + 2)}{2(z^2 + 1)}t^2 + \frac{z}{z^2 + 1}t + \frac{3}{2(z^2 + 1)}.$$

Continue the reduction procedure of Section 3.2 one obtains

$$u' + u^2 = Bu + C,$$

where

$$B = -\frac{z^3}{(z^2 + 1)(z^2 + 2)} \quad \text{and} \quad C = \frac{3(z^2 + 2)}{4(z^2 + 1)^2}.$$

And finally one obtains the following classical Riccati equation

$$v' + v^2 = -6(z^2 + 2)^{-2},$$

which has a rational general solution as follows

$$v(z) = -\frac{z}{z^2 + 2} + \frac{1}{z - \lambda} + \frac{1}{z + 2/\lambda}$$

with λ the arbitrary constant. One finally has by substitutions the following solution of equation $F(z, w, w') = 0$:

$$w(z) = \frac{z^2 \lambda^2 - 2z\lambda^3 + 4z\lambda + 4 + \lambda^4 - 3\lambda^2}{(z\lambda + 2 - \lambda^2)^2 z}.$$

5. CONCLUSION

In this paper, we present an algebraic geometry approach to the study of first order algebraic differential equations. The necessary and sufficient conditions of the rational general solutions of the Riccati equations are studied in [6]. In [7], we prove that the degree of a rational general solution of a first order algebraic differential equation of degree d is less than or equal to d . The algebraic geometry approach is also used to obtain bound of the number of the rational solutions of a first order algebraic differential equation with algebraic genus greater than one. In [8], the algebraic general solutions of first order algebraic differential equations were studied by using of the birational transformations of algebraic curves, and an algorithm was presented to get an algebraic general solution of first order algebraic differential equations without movable critical point if the algebraic general solution exists. It is interesting to present an effective algorithm to find the rational general solutions or the rational solutions of a first order algebraic differential equation.

REFERENCES

- [1] S. S. Abhyankar and C. L. Bajaj, Automatic parameterization of rational curves and surfaces. III. Algebraic plane curves. *Comput. Aided Geom. Design*, **5** (1988), 309–321.
- [2] C. Briot and T. Bouquet, Intégrations des équations différentielles au moyen des fonctions elliptiques, *Journal de l'Ecole Polytechnique*, t. XXI.
- [3] M. Bronstein, Linear ordinary differential equations: breaking through the order 2 barrier, *Proceedings of ISSAC 1992*, ACM Press, 42–48.
- [4] J. G. Campbell, A criterion for the polynomial solutions of a certain Riccati equation, *American Math. Monthly*, **61** (1952), 388–389.
- [5] M. M. Carnicer, The Poincaré problem in the nondicritical case. *Ann. of Math. (2)* **140** (1994), no. 2, 289–294.
- [6] G. Chen and Y. Ma, *A computer algebra approach to first order algebraic differential equations*. Constructive and Invariant Methods In Algebraic and Differential Equations — The Sixth International Workshop on Mathematics Mechanization, Shanghai, China, May 19–21, 2004.
- [7] G. Chen and Y. Ma, *Rational solutions of algebraic differential equations*. The Third International Congress of Chinese Mathematicians, Hong Kong, December 17–22, 2004.
- [8] G. Chen and Y. Ma, *Algebraic solutions of a first order algebraic differential equation*, Preprint, Laboratory of P. Painlevé, 2004.
- [9] A. Eh. Eremenko, Rational solutions of first-order differential equations. *Ann. Acad. Sci. Fenn.*, **23** (1998), 181–190.
- [10] R. Feng and X. Gao, Rational general solutions of algebraic ordinary differential equations, *Proceeding of ISSAC 2004*, ACM Press, 155–162.
- [11] L. Fuchs, Ueber Differentialgleichungen deren integrale feste verzweigungspunkte besitzen, *Sitzungsberichte Acad. Berlin*, 1884.

- [12] V. V. Golubev, Lectures on the analytic theory of differential equations. 2nd ed. Moscow–Leningrad: gos. Izd. Tekh. Teor. Lit. **436** p. (1950). (Russian, Chinese translation and German translation are available).
- [13] E. L. Ince, *Ordinary Differential Equations*, London, Longmans, Green and Co. Ltd., 1927.
- [14] J. J. Kovacic, An algorithm for solving second order linear homogeneous differential equations, *J. Symbolic Comp.*, **2** (1986), 3–43.
- [15] Z. M. Li and F. Schwarz, Rational solutions of Riccati-like partial differential equations. *J. Symbolic Comp.*, **31** (2001), no. 6, 691–716.
- [16] Y. Ma and X. Gao, Polynomial solutions of algebraic differential equations. Lecture Notes Ser. Comput., **9**, World Sci. Publishing, River Edge, 2001, 92–101.
- [17] M. Matsuda, *First Order Algebraic Differential Equations — A Differential Algebraic Approach*, Lecture Notes in Math. **804**, Springer-Verlag, Berlin, 1980.
- [18] P. Painlevé, Leçons sur la théorie analytique des équations différentielles, 1896.
- [19] É. Picard, *Traité D’Analyse*, Tome III, 1908.
- [20] H. Poincaré, Sur un théorème de M. Fuchs, *Acta Math.*, **7** (1885), 1–32. Also in *Oeuvres de Henri Poincaré*, Vol. 1, Gauthiers-Villars et Cie, Paris, 1928.
- [21] R. H. Risch, The problem of integration in finite terms, *Trans. Amer. Math. Soc.*, **139** (1969), 167–189.
- [22] T. W. Sederberg, D. C. Anderson and R. N. Goldman, Implicit representation of parametric curves and surfaces, *Computer Vision, Graphics, and Image Processing*, **28** (1984), 72–84.
- [23] T. W. Sederberg, J. Zheng, Algebraic methods for computer aided geometric design. *Handbook of computer aided geometric design*, 363–387, North-Holland, Amsterdam, 2002.
- [24] J. R. Sendra and F. Winkler, Symbolic parametrization of curves. *J. Symbolic Comp.*, **12** (1991), 607–631.
- [25] J. R. Sendra and F. Winkler, Parametrization of algebraic curves over optimal field extension. *J. Symbolic Comp.*, **23** (1997), 191–207.
- [26] M. F. Singer, Liouvillian solutions of linear differential equations with Liouvillian coefficients. *J. Symbolic Comp.*, **11** (1991), 251–273.
- [27] M. van Hoeij, Rational parametrizations of algebraic curves using a canonical divisor. *J. Symbolic Comp.*, **23** (1997), 209–227.
- [28] R. J. Walker, *Algebraic Curves*, Princeton University Press, Princeton, 1950.
- [29] W. T. Wu, Mechanical Derivation of Newton’s Gravitational Laws from Kepler’s Laws, MM Research Preprints, No. **1** (1987), 53–61.
- [30] W. T. Wu, On the Foundation of Algebraic Differential Geometry, MM Research Preprints, No. **3** (1989), 1–26.
- [31] W. T. Wu, On algebrico-differential equations-solving, MM Research Preprints, No. **22**, December 2003, 1–13.
- [32] W. Yuan, A note on the Riccati differential equation, *J. Math. Anal. Appl.*, **277** (2003), 367–374.

Acknowledgment. The authors are grateful to the anonymous referee for the valuable comments and suggestions, which improved the manuscript, and for pointing out some inaccuracies.

LABORATOIRE P. PAINLEVÉ, UMR CNRS 8524, UFR DE MATHÉMATIQUES, UNIVERSITÉ DE LILLE 1, 59655 VILLENEUVE D’ASCQ, FRANCE
E-mail address: Guoting.Chen@math.univ-lille1.fr

KEY LABORATORY OF MATHEMATICS MECHANIZATION, CHINESE ACADEMY OF SCIENCES, BEIJING 100080, P. R. CHINA
E-mail address: yjma@mmrc.iss.ac.cn